



Laccase C

Benzenediol: oxygen oxidoreductase, E.C. 1.10.3.2

Description: Enzyme preparation which oxidised monovalent and polyvalent phenolic compounds. It catalyses reactions to products similar like humic substances under less oxygenic conditions. Basically, Laccase oxidises ortho- and para-diphenols into the corresponding quinones and phenoxy radicals which polymerises spontaneously and are precipitated in the solution.

Catalysed reaction:



Origin: *Trametes spec.*

Application:

- Precipitation of phenolic substances
- Enzymatic browning of food (cacao, coffee)
- Glueing of flake boards
- Modification of elasticity and consistency of pastes, gums, dispersion media, phenolic resins
- Production of microbicides
- Analysis of phenols

Activity: > 500 U/ g
(Substrate: Syringaldazin)

Specific activity: > 75 U/ mg Protein

Parameter:	pH	Optimum: 5.0, effective in the range of pH 3.0 - 7.5
	Temperature	Optimum: 60°C, effective in the range of 20 - 70°C
Article-no.:	2020	
Form of delivery:	Light brown powder/ lyophilizate	
Stability:	Stable at 4 – 6 °C, temperatures down to -20°C are possible	
Storage:	at 4 – 6°C, storage down to – 20°C is possible	
Literature:	[1] Wood, D.A., (1979), J. Gen. Microbiol., <u>117</u> , 327-338	