Cutinase 2P

EC 3.1.1.74

Beschreibung: Enzympräparat zur Hydrolyse von Cutin und anderen Estern. Cutin (Polyester aus Hydroxy- und Hydroxy-Epoxyfettsäure) wird in die entsprechenden Monomere gespalten.

Katalysierte Reaktion:

Cutin + *H_2O* → *2 Cutin Monomere*

Herkunft: *Arxula adenivorans*

Aktivität: > 60 000 U/ g (pH 7,0; 37°C; Glycerintributyrat als Substrat)

Spezifische Aktivität: > 3 000 U/ mg Protein

Anwendung: Spaltung von Cutin; organische Synthese (s. Tab. 1)

Reaktionsparameter:			
	pH–Wert	Optimum: 4 - 6	aktiv im Bereich pH 3,5 - 8
	Temperatur	Optimum: 20 - 40°C	aktiv im Bereich 5 - 60°C

ASA Spezialenzyme GmbH
Am Exer 19c
38302 Wolfenbüttel
Germany

Tel.: +49 (0)5331 8825-30
Fax: +49 (0)5331 8825-32

E-Mail: service@asa-enzyme.de
Internet: www.asa-enzyme.de



Abbildung 1: pH-Abhängigkeit



Abbildung 2: pH-Stabilität



Abbildung. 3: Temperatur-Abhängigkeit der Cutinase 2P

Artikel-Nr.:	2465
Lieferform:	Teilgereinigtes Lyophilisat
Lagerung:	- 20°C

Tab. 1: Einsatz von Cutinasen in der Biokatalyse

(nach Carvalho et al., 1998b)

Reaktion	Substrat	Enzympräparation/Reaktionsansatz	Referenz.
Hydrolyse	Triolein	Reverse Micellen AOT/Isoktan	Melo et al., 1995b
		Triolein/Wasser	Flipsen et al., 1996
	Tricaprylin	Immobilisierung an Zeolithen	Gonçalves et al., 1996a
		Calciumalginat	Gonçalves et al., 1995
		Kovalente Bindung an porösem Silicat	Gonçalves et al., 1996b
	p-nitrophenyl valerat	Micellen mit SDS/Triton X100	Pocalyko and Tallman, 1998
	p-nitrophenyl palmitat	Immobilizierung an Dextran und Silica-Derivate	Gonçalves et al., 1998a
	Methyl-,ethyl-, propylpropionat	Gas/Festphasensystem	Lamare et al., 1997
Estersynthese	Ölsäure+Hexanol	Reverse Micellen AOT/Isoktan	Sebastiãoet al., 1993, Sebas tião et al., 1992
	Caprylsäure + Butanol	Organische Lösungsmittel	Sarazin et al., 1992, Sarazin et al., 1995
	Caprylsäure + Butanol	Organische Lösungsmittel	Sarazin et al., 1992, Sarazin et al., 1995
	Buttersäure + 2-Butanol	Phosphatidylcholin/ Isoktan, Reverse Micellen	Pinto-Sousa et al., 1994
	Ölsäure + Glycerin	Organische Lösungsmittel	Melo et al., 1995a
	Hexansäure + Hexanol	CTAB, Reverse Micellen	Cunnah et al., 1996
		Immobilisierung an Accurel EP 100	Sereti et al., 1997
	Buttersäure + Hexanol	Immobilisierung an Accurel EP 100	Sjursnes et al., 1998
	Laurinsäure + Pentanol	Reverse Micellen AOT/Isoktan	Papadimitriou et al.,1996
Umesterung	Methylpropionat+Propanol	Gas/Festphasensystem	Lamare and Legoy, 1995, Lamare et al., 1997
	Butylacetat+Hexanol	Reverse Micellen AOT/Isoktan	Carvalho et.al 1997a, Carvalho et al., 1998a
		Reverse Micellen CTAB/Isoktan	Cunnah et al., 1996
		Immobilisierung an Zeolithen	Serralha et al., 1998

Literatur

Carvalho, C.M.L., Serralheiro, M.L.M., Cabral, J.M.S. and Aires-Barros, M. R. (1997a). Application of factorial design to the study of transesterification reactions using cutinase in AOT-reversed micelles. Enzyme and Microbiological Technology 21: 117-123.

Carvalho, C.M.L., Cabral, J.M.S. and Aires-Barros, M. (1998a) Kinetics and modelling of transesterification reactions catalysed by cutinase in AOT reversed micelles. Journal of Molecular Catalalysis B- Enzymatic 5: 361-365.

Carvalho, C. M. L., Aires-Barros, M. R. and Cabral , J. M. S. (1998b). Cutinase structure, function and biocatalytic applications EJB Electronic Journal of Biotechnology, 1: 160-173.

Cunnah, P.J., Aires-Barros, M.R. and Cabral, J.M.S. (1996). Esterification and transesterification catalysed by cutinase in reverse micelles of CTAB for the synthesis of short chain esters. Biocatalysis and Biotransformation 14:125-146.

Flipsen, J.A.C., van der Hijden, H.T.W.M., Egmond, M.R. and Verheij, H.M. (1996) Action of cutinase at the trioleinwater interface. Characterisation of interfacial effects during lipid hydrolysis using the oil-drop tensiometer as a tool to lipase kinetics. Chemistry and Physics of Lipids 84:105-115.

Gonçalves, A.P.V., Cabral, J.M.S. and Aires-Barros, M.R. (1995) Immobilization of a recombinant cutinase by entrapment and by covalent binding. Applied Biochemistry and Biotechnology 60:217-228.

Gonçalves, A.P.V., Lopes, J.M., Lemos, F., Ramôa Ribeiro, F., Prazeres, D.M.F., Cabral, J.M.S. and Aires-Barros, M.R.(1996a) Zeolites as supports for enzymatic hydrolysis reactions. Comparative study of several zeolites. Journal of Molecular Catalysis B- Enzymatic 1:53-60.

Gonçalves, A.M., Schacht, E., Cabral, J.M.S. and Gil, M.H. (1996b) Stability studies of a recombinant cutinase immobilised on an inorganic support. In: Proceedings of III Congreso Ibérico de Biotecnologia, F.F. Polanco, P.G. Encina, G.G. Benito and M.M. P. Miranda, Universidad de Valladolid, pp. 551-552.

Gonçalves, A.M., Schacht, E., Matthijs, G., Aires-Barros, M.R., Cabral, J.M.S., Gil, M.H. (1998a) Stability studies of a recombinant cutinase immobilized to dextran and derivatized silica supports. Enzyme and Microbial Technology. In press.

Lamare, S. and Legoy, M.D. (1995) Working at controlled water activity in a continuous process: the gas/solid system as a solution. Biotechnology and Bioengineering 45:387-397.

Lamare, S., Lortie, R. and Legoy, M.D. (1997) Kinetic studies of *Fusarium solani pisi* cutinase used in a gas/solid system: transesterification and hydrolysis reactions. Biotechnology and Bioengineering 56:1-7.

Melo, E.P., Ivanova, M.G., Aires-Barros, M.R., Cabral. J.M.S. and Verger, R. (1995a) Glyceride synthesis catalyzed by cutinase using the monomolecular film technique. Biochemistry 34:1615-1621.

Melo, E.P., Aires-Barros, M.R., Cabral, and J.M.S. (1995b) Triglyceride hydrolysis and stability of a recombinant cutinase from *Fusarium solani* in AOT-*iso*-octane reversed micelles. Appl. Biochem. Biotechnol. 50:45-56.

Papadimitriou, V., Xenakis, A., Cazianis, C.T., Stamatis, H., Egmond, M. and Kolisis, F.N. (1996) EPR studies of cutinase in microemulsions Annals NewYork Academy of Sciences 799:275-280.

Pinto-Sousa, A.M., Cabral, J.M.S. and Aires-Barros, M.R. (1994) Ester synthesis by a recombinant cutinase in reversed micelles of a natural phospholipid. Biocatalysis 9: 169-179.

Pocalyko, D.J. and Tallman, M. (1998) Effects of amphipaths on the activity and stability of *Fusarium solani pisi* cutinase. Enzyme and Microbial Technology 22: 647- 651.

Sarazin, C., Goethals, G., Séguin, J.P., Legoy, M.D. and Barbotin, J.N. (1992) Usefulness of NMR methods for assaying cutinase catalysed synthesis of ester in organic media. In: Biocatalysis in Non-Conventional Media, J. Tramper, M.H. Vermue, H.H. Beeftink and U. von Stockar. Elsevier Science Publishers B.V., 1992, pp. 23-29.

Sarazin, C., Ergan, F., Séguin, J.P., Goethals, G., Legoy M.D. and Barbotin J.N. (1995) NMR on-line monitoring of esterification catalyzed by cutinase. Biotechnology and Bioengineering 51:636-644.

Sebastião, M. J., Cabral, J.M.S., and Aires-Barros, M.R. (1992) Synthesis of fatty acid esters by a recombinant cutinase in reversed micelles. In: Biocatalysis in Non- Conventional Media, J. Tramper, M.H. Vermue, H.H. Beeftink and U. von Stockar. Elsevier Science Publishers B.V., 1992, pp. 719-724.

Sebastião, M. J., Cabral, J.M.S., and Aires-Barros, M.R. (1993) Synthesis of fatty acid esters by a recombinant cutinase in reversed micelles. Biotechnology and Bioengineering 42:326-332.

Serralha, F.N., Lopes, J.M., Lemos, F., Prazeres, D.M.F., Aires-Barros, M.R. Cabral, J.M.S., and Ribeiro, F. R. (1998) Zeolites as supports for an enzymatic alcoholysis reaction . Journal of Molecular Catalysis B- Enzymatic 4:303-311.

Sereti, V., Stamatis, H. and Kolisis, F.N. (1997) Improved stability and reactivity of *Fusarium solani* cutinase in supercritical CO2. Biotechnology Techniques 11:661-665.

Sjursnes, B., Valente, C. and Halling, P. (1998) Interactions between water and medium effects on enzymes in organic media: kinetics of cutinase catalysed esterification with mutual solvation of reactants. Biotechnology and Bioengineering In press.